

## TRANSFORMATION OF RANDOM VARIABLES

Let  $(X, Y)$  be a continuous two dimensional random variables with JPDF  $f_{XY}(x, y)$ . Transform  $X$  and  $Y$  to new random variables  $U = h(x, y), V = g(x, y)$ .

Then the joint PDF of  $(U, V)$  is given by  $f_{UV}(u, v) = |J| f_{XY}(x, y)$

$$\text{where } J = \frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$$

### PROCEDURE TO FIND THE MARGINAL PDF OF $U$ & $V$

(1) Take  $u$  as the RV to which the PDF to be computed and take  $v = y$ . (if not given)

(2) Express  $x$  and  $y$  in terms of  $u$  and  $v$ .

(3) Find  $J = \frac{\partial(x,y)}{\partial(u,v)} = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix}$

(4) Write the JPDF of  $(U, V), f_{UV}(u, v) = |J| f_{XY}(x, y)$

(5) Substitute the values of  $J, x$  and  $y$ .

(6) Find the range of  $u$  and  $v$  using the range of  $x$  and  $y$ .

(7) The PDF of  $U$  is  $f_U(u) = \int_{v=-\infty}^{v=\infty} f_{uv}(u, v) dv$

(8) The PDF of  $V$  is  $f_V(v) = \int_{u=-\infty}^{u=\infty} f_{uv}(u, v) du$

**Problem based on transformation of random variables**

**1. If the JPDF  $f(x, y)$  is given by  $f_{XY}(x, y) = x + y; 0 \leq x, y \leq 1$ , find PDF of  $U = XY$ .**

**Solution:**

Given  $(X, Y)$  is a continuous 2D RV defined in  $0 < x < 1$  and  $0 < y < 1$ .

Also Given  $f_{xy}(x, y) = x + y; 0 \leq x, y \leq 1$

we have to find the PDF of  $u = xy \dots \dots \dots (1)$

let  $v = y \Rightarrow y = v$ .

(1)  $\Rightarrow u = xv \Rightarrow x = \frac{u}{v}$

$\therefore x = \frac{u}{v} \quad y = v$

$\frac{\partial z}{\partial u} = \frac{1}{v}; \frac{\partial x}{\partial v} = \frac{-u}{v^2}; \frac{\partial y}{\partial u} = 0; \frac{\partial y}{\partial v} = 1$

$J = \begin{vmatrix} \frac{1}{v} & \frac{-u}{v^2} \\ 0 & 1 \end{vmatrix} = \frac{1}{v}$

$J = \frac{1}{v}$

The JPDF of  $(U, V) f_{uv}(u, v) = |J| f_{xy}(x, y)$

$$= \left| \frac{1}{v} \right| (x + y) = \frac{1}{v} \left( \frac{u}{v} + v \right)$$

$$= \frac{u}{v^2} + 1$$

$$f_{uv}(u, v) = \frac{u}{v^2} + 1$$

**To find the range for  $u$  and  $v$ :**

We have  $0 \leq x \leq 1 \Rightarrow 0 \leq \frac{u}{v} \leq 1$

i.e  $0 \leq u \leq v$

Also  $0 \leq y \leq 1 \Rightarrow 0 \leq v \leq 1$

On combining the two limits, we get  $0 \leq u \leq v \leq 1$

$$\therefore f_{uv}(u, v) = \frac{u}{v^2} + 1, 0 \leq u \leq v \leq 1$$

**PDF of  $U$  is given by**

$$f_U(u) = \int_{v=u}^{v=1} f_{uv}(u, v) dv \quad 0 \leq u \leq v < 1$$

$$= \int_u^1 \left( \frac{u}{v^2} + 1 \right) dv$$

$$= \int_u^1 (uv^{-2} + 1) dv$$

$$= \left[ \frac{uv^{-1}}{-1} + v \right]_u^1$$

$$= \left( \frac{u}{-1} + 1 \right) + 1 - u$$

$$= -u + 1 + 1 - u$$

$$= 2 - 2u$$

$$f_U(u) = 2(1 - u) \quad 0 < u < 1$$

2. Let  $(X, Y)$  be a continuous two dimensional random. with JPDF

$$f(x, y) = 4xye^{-(x^2+y^2)} \quad x > 0, y > 0. \text{ Find the PDF of } \sqrt{X^2 + Y^2}$$

**Solution:**

Given  $(X, Y)$  is a continuous 2D RV defined in  $0 < x < \infty$  and  $0 < y < \infty$

Given  $f(x, y) = 4xye^{-(x^2+y^2)}$ ,  $0 < x < \infty, 0 < y < \infty$

let  $u = \sqrt{x^2 + y^2}$  ..... (1)      Take  $v = y \Rightarrow y = v$

$$(1) \Rightarrow u^2 = x^2 + y^2$$

$$u^2 = x^2 + y^2 \quad y = v$$

$$x^2 = u^2 - v^2 \Rightarrow x = \sqrt{u^2 - v^2}$$

$$x\sqrt{u^2 - v^2}, y = v$$

$$\frac{\partial x}{\partial u} = \frac{1}{2} \frac{1}{\sqrt{u^2 - v^2}} (2u) = \frac{u}{\sqrt{u^2 - v^2}}; \frac{\partial y}{\partial u} = 0$$

$$\frac{\partial x}{\partial v} = \frac{1}{2} \frac{1}{\sqrt{u^2 - v^2}} (-2v) = \frac{-v}{\sqrt{u^2 - v^2}}; \frac{\partial y}{\partial v} = 1 = 1$$

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} \frac{u}{\sqrt{u^2 - v^2}} & \frac{-v}{\sqrt{u^2 - v^2}} \\ 0 & 1 \end{vmatrix}$$

$$J = \frac{u}{\sqrt{u^2 - v^2}}$$

PDF of  $(U, V)$  is  $f_{UV}(u, v) = |J| f_{XY}(x, y)$

$$= \frac{u}{\sqrt{u^2 - v^2}} 4xy e^{-(x^2 + y^2)}$$

$$= \frac{u}{\sqrt{u^2 - v^2}} 4\sqrt{u^2 - v^2}(v) e^{-u^2}$$

$$f_{UV}(u, v) = 4uve^{-u^2}$$

**To find the range for  $u$  and  $v$ :**

We have  $x > 0$

We have  $y > 0$

$$\sqrt{u^2 - v^2} > 0$$

$$v > 0$$

$$u^2 - v^2 > 0$$

$$\Rightarrow 0 < v < \infty$$

$$u^2 > v^2 \Rightarrow u > v$$

$$\Rightarrow v < u$$

On combining the two limits, we get  $0 < v < u < \infty$

$$f_{UV}(u, v) = 4uve^{-u^2}, 0 < v < u < \infty$$

**PDF of  $U$  is given by**

$$\begin{aligned} F_U(u) &= \int_0^u f_{UV}(u, v) dv \\ &= \int_0^u 4uve^{-u^2} dv \\ &= 4ue^{-u^2} \int_0^u v dv \\ &= 4ue^{-u^2} \left[ \frac{v^2}{2} \right]_0^u \\ &= 2u^3 e^{-u^2} \quad 0 < u < \infty \end{aligned}$$

**3. The JPDF to two diamensional random variables  $X$  and  $Y$  is given by,**

$$(x, y) = e^{-(x+y)}, x > 0, y > 0. \text{ Find the PDF of } \frac{X+Y}{2}$$

**Solution:**

Given  $(X, Y)$  is a continuous 2DRV defined in  $0 < x < \infty$  and  $0 < y < \infty$ .

Also given  $f(x, y) = e^{-(x+y)}$ ;  $0 < x < \infty, 0 < y < \infty$

let  $u = \frac{x+y}{2}$  ..... (1). Take  $v = y \Rightarrow y = v$

$$(1) \Rightarrow u = \frac{1}{2}(x + v)$$

$$2u = x + vx = 2u - v$$

$$\Rightarrow x = 2u - v; y = v \frac{\partial x}{\partial u} = 2 \frac{\partial x}{\partial v} = -1; \frac{\partial y}{\partial u} = 0; \frac{\partial y}{\partial v} = 1$$

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 2 & -1 \\ 0 & 1 \end{vmatrix} = 2$$

the PDF of  $(U, V)$  is  $f_{uv}(u, v) = |J|f_{xy}(x, y)$   
 $= 2e^{-(x+y)}$

$$= 2e^{-(2u-v+v)}$$

$$= 2e^{-2u}$$

**To find range for  $u$  and  $v$  :**

We have  $x > 0 \Rightarrow 2u - v > 0$

i. e.,  $2u > v \Rightarrow v < 2u$

Also  $y > 0 \Rightarrow v > 0$

$$\therefore v < 2u; v > 0$$

$$0 < v < 2u < \infty$$

On combining the two limits, we get  $0 < v < 2u < \infty$

$$\therefore f_{UV}(u, v) = 2e^{-2u}, 0 < v < 2u < \infty$$

**The PDF of  $U$  is**

$$f_U(u) = \int_{v=0}^{v=2u} f_{uv}(u, v) dv$$

$$= \int_0^{2u} 2e^{-2u} dv$$

$$= 2e^{-2u} \int_0^{2u} dv$$

$$= 2e^{-2u} [v]_0^{2u}$$

$$= 2e^{-2u} (2u)$$

$$f_u(u) = 4ue^{-2u}; u > 0$$

**Unit step function:**

$$u(x) = 1 \text{ for } x > 0$$

$$u(x) = 0 \text{ for } x < 0$$

**1. If  $X$  and  $Y$  are two independent random variables each normally distributed with mean = 0 and variance  $\sigma^2$ , find the density function of**

$$R = \sqrt{X^2 + Y^2} \text{ and } \phi = \tan^{-1} \left( \frac{Y}{X} \right)$$



**Solution:**

Given that  $X$  follows  $N(0, \sigma)$ .

$$\therefore f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2\sigma^2}x^2}; -\infty < x < \infty$$

Also  $Y$  follows  $N(0, \sigma)$ .

$$\therefore f_Y(y) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2\sigma^2}y^2}; -\infty < y < \infty$$

Since  $X$  and  $Y$  are independent,  $f_{XY}(x, y) = f_X(x)f_Y(y)$

$$= \frac{1}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2}(x^2+y^2)}; -\infty < x < \infty, -\infty < y < \infty$$

We have  $r = \sqrt{x^2 + y^2}; \theta = \tan^{-1}\left(\frac{y}{x}\right)$

$$\Rightarrow x = r\cos\theta, y = r\sin\theta, J = r$$

JPDF of  $(R, \phi)$  is  $f_{R\phi}(r, \theta) = |J| f_{XY}(x, y)$

$$= r \frac{1}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2}(x^2+y^2)}$$

$$= \frac{r}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2}r^2}$$

**To find the range for  $r$  and  $\theta$  :**

We have  $-\infty < x < \infty, -\infty < y < \infty$  t.e entire  $XY$  plane.

The entire  $XY$  plane is transformed into  $x = r\cos\theta, y = r\sin\theta$

i.e the entire  $XY$  plane is transformed into  $x^2 + y^2 = r^2$  ( a circle of infin

radius)

Whole region is transformed into a circle of infinite radius.

$$\therefore 0 \leq r < \infty, 0 \leq \theta \leq 2\pi$$

$$\therefore f_{R\phi}(r, \theta) = \frac{r}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2} r^2} \quad 0 \leq r < \infty, 0 \leq \theta \leq 2\pi$$

**The PDF of R is**

$$\begin{aligned} f_R(r) &= \int_{r=0}^{\infty} f_{r\theta}(r, \theta) d\theta \\ &= \int_0^{2\pi} \frac{r}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2} r^2} d\theta \\ &= \frac{r}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2} r^2} \int_0^{2\pi} d\theta \\ &= \frac{r}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2} r^2} [\theta]_0^{2\pi} \end{aligned}$$

$$f_R(r) = \frac{r}{\sigma^2} e^{-\frac{1}{2\sigma^2} r^2}; 0 \leq r < \infty$$

**The PDF of  $\phi$  is**

$$\begin{aligned} f_\phi(\theta) &= \int_{r=0}^{\infty} f_{r\theta}(r, \theta) dr \\ &= \int_0^{\infty} \frac{r}{\sigma^2 2\pi} e^{-\frac{1}{2\sigma^2} r^2} dr \\ &= \frac{1}{\sigma^2 2\pi} \int_0^{\infty} r e^{-\frac{1}{2\sigma^2} r^2} dr \end{aligned}$$

$$\text{Put } \frac{1}{2\sigma^2} r^2 = t$$

$$\frac{1}{2\sigma^2} 2rdr = dt$$

$$rdr = \sigma^2 dt$$

There is no change on the limits

$$\begin{aligned} f_0(\theta) &= \frac{1}{\sigma^2 2\pi} \int_0^{\infty} e^{-t} \sigma^2 dt \\ &= \frac{1}{2\pi} \left[ \frac{e^{-t}}{-1} \right]_0^{\infty} \\ &= \frac{1}{2\pi} (0 + 1) \end{aligned}$$

$$f_\phi(\theta) = \frac{1}{2\pi} \quad 0 \leq \theta \leq 2\pi$$

**2. The random variables  $X$  and  $Y$  each follows an exponent distribution with parameter 1 and are independent. Find the PDF of  $U = X - 1$**

**Solution:**

Given  $X$  and  $Y$  follows exponential distribution with parameter with  $\lambda = 1$

$$\therefore f_x(x) = \lambda e^{-\lambda x}; x > 0$$

$$= e^{-x}$$

$$f_y(y) = e^{-y}; y > 0$$

Since  $X$  and  $y$  are independent,

$$f_{XY}(x, y) = f_x(x)f_y(y)$$

$$= e^{-x}e^{-y}$$

$$= e^{-(x+y)}$$

let  $u = x - y$  .....(1) Take  $v = y \Rightarrow y = v$

$$(1) \Rightarrow u = x - v \Rightarrow x = u + v$$

$$x = u + v ; y = v$$

$$J = \begin{vmatrix} \frac{\partial x}{\partial u} & \frac{\partial x}{\partial v} \\ \frac{\partial y}{\partial u} & \frac{\partial y}{\partial v} \end{vmatrix} = \begin{vmatrix} 1 & 1 \\ 0 & 1 \end{vmatrix} = 1$$

The JPFD of  $(U, V)$  is  $f_{uv}(u, v) = |J|f_{XY}(x, y)$

$$= (1)e^{-(x+y)}$$

$$= e^{-(u+v+v)}$$

$$= e^{-(u+2v)}$$

**To find the range for  $u$  and  $v$  :**

We have  $x > 0 \Rightarrow u + v > 0 \Rightarrow u > -v$

$$\text{for } y > 0 \Rightarrow v > 0$$

$$\therefore f_{uv}(u, v) = e^{-(u+2v)}u > -v, v > 0$$

**The PDF of  $U$  is**

$$f_u(u) = \int f(u, v)dv$$

Since there are two slopes, the region is divided into two sub regions  $R_1$  and  $R_2$

In  $R_1$ :

$$\text{At } P_1, v = -u; \text{ At } Q_1, v = \infty$$

In  $R_2$ :

$$\text{At } P_2, v = 0; \text{ At } Q_2, v = \infty$$

$$\text{In } R_1 : f_U(u) = \int_{-u}^{\infty} f(u, v)dv$$

$$= \int_{-u}^{\infty} e^{-(u+2v)}dv$$

$$= \int_{-u}^{\infty} e^{-u} e^{-2v} dv$$

$$= e^{-u} \int_{-u}^{\infty} e^{-2v} dv$$

$$= e^{-u} \left[ \frac{e^{-2v}}{-2} \right]_{-u}^{\infty}$$

$$= e^{-u} \left[ 0 - \frac{e^{2u}}{-2} \right]$$

$$= \frac{e^u}{2}; u < 0$$

In  $R_2$

$$f_U(u) = \int_{v=0}^{\infty} e^{-u} f(u, v) dv$$

$$= \int_0^{\infty} e^{-(u+2v)} dv$$

$$= \int_0^{\infty} e^{-u} e^{-2v} dv$$

$$= \int_0^{\infty} e^{-u} \left[ 0 - \frac{1}{-2} \right]$$

$$= \frac{e^{-u}}{2}; u > 0$$

$$= e^{-u} \int_0^{\infty} e^{-2v} dv$$

$$= e^{-u} \left[ \frac{e^{-2v}}{-2} \right]_0^{\infty}$$

$$f_U(u) = \begin{cases} \frac{e^u}{2} & u < 0 \\ \frac{e^{-u}}{2} & u > 0 \end{cases}$$

OBSERVE OPTIMIZE OUTSPREAD