

## MATRIX OF LINEAR TRANSFORMATION WITH STANDARD BASES

1. Find the matrix of the linear transformation  $T: R^2 \rightarrow R^2$  given by  $T(a, b) = (2a - 3b, a + b)$  relative to the basis (i)  $\{(1, 0), (0, 1)\}$   
(ii)  $\{(2, 3), (1, 2)\}$

Solution

Given,  $T(a, b) = (2a - 3b, a + b)$

(i) The standard bases of  $R^2$  is  $\beta = \gamma = \{(1, 0), (0, 1)\}$

Given,  $T(a, b) = (2a - 3b, a + b)$

$\therefore$  the matrix of the linear transmission is  $[T]_{\beta}^{\gamma} = \begin{bmatrix} 2 & -3 \\ 1 & 1 \end{bmatrix}$

(ii) the basis is  $\beta = \{(2, 3), (1, 2)\}$

$$v_1 = (2, 3), v_2 = (1, 2)$$

$$T(a, b) = (2a - 3b, a + b)$$

$$T(v_1) = T(2, 3)$$

$$= (2(2) - 3(3), 2 + 3)$$

$$= (-5, 5)$$

The first column of the matrix of  $T$  is  $\begin{bmatrix} -5 \\ 5 \end{bmatrix}$

$$T(v_2) = T(1, 2)$$

$$= (2(1) - 3(2), 1 + 2)$$

$$= (-4, 3)$$

The second column of the matrix of  $T$  is  $\begin{bmatrix} -4 \\ 3 \end{bmatrix}$

Matrix of  $T$  is  $\begin{bmatrix} -5 & -4 \\ 5 & 3 \end{bmatrix}$

2. Let  $T: V_2(\mathbb{R}) \rightarrow V_3(\mathbb{R})$  and  $U: V_2(\mathbb{R}) \rightarrow V_3(\mathbb{R})$  be the linear transformations respectively defined by  $T(a_1, a_2) = (a_1 + 3a_2, 0, 2a_1 - 4a_2)$  and  $U(a_1, a_2) = (a_1 - a_2, 2a_1, 3a_1 + 2a_2)$ . Let  $\beta$  and  $\gamma$  be the standard bases of  $V_2(\mathbb{R})$  and  $V_3(\mathbb{R})$  respectively. Verify  $[T + U]_{\beta}^{\gamma} = [T]_{\beta}^{\gamma} + [U]_{\beta}^{\gamma}$

Solution:

Given,  $T(a_1, a_2) = (a_1 + 3a_2, 0, 2a_1 - 4a_2)$

Since  $\beta$  and  $\gamma$  be the standard bases of  $V_2(\mathbb{R})$  and  $V_3(\mathbb{R})$

the matrix corresponds to  $\beta = [T]_{\beta}^{\gamma} = \begin{bmatrix} 1 & 3 \\ 0 & 0 \\ 2 & -4 \end{bmatrix} \dots (1)$

Given,  $U(a_1, a_2) = (a_1 - a_2, 2a_1, 3a_1 + 2a_2)$ .

Since  $\beta$  and  $\gamma$  be the standard bases of  $V_2(\mathbb{R})$  and  $V_3(\mathbb{R})$

the matrix corresponds to  $\beta = [U]_{\beta}^{\gamma} = \begin{bmatrix} 1 & -1 \\ 2 & 0 \\ 3 & 2 \end{bmatrix} \dots (2)$

$$(1) + (2) \Rightarrow [T]_{\beta}^{\gamma} + [U]_{\beta}^{\gamma} = \begin{bmatrix} 2 & 2 \\ 2 & 0 \\ 5 & -2 \end{bmatrix} \dots (3)$$

$$\begin{aligned} (T + U)(a_1, a_2) &= (a_1 + 3a_2, 0, 2a_1 - 4a_2) + (a_1 - 3a_2, 2a_1, 3a_1 + 2a_2) \\ &= (a_1 + 3a_2 + a_1 - 3a_2, 0 + 2a_1, 2a_1 - 4a_2 + 3a_1 + 2a_2) \\ &= (2a_1 + 2a_2, 2a_1, 5a_1 - 2a_2) \end{aligned}$$

Since  $\beta$  and  $\gamma$  be the standard bases of  $V_2(R)$  and  $V_3(R)$

the matrix corresponds to  $\beta = [T + U]_{\beta}^{\gamma} = \begin{bmatrix} 2 & 2 \\ 2 & 0 \\ 5 & -2 \end{bmatrix} \dots (4)$

$$\text{From (3) and (4)} \Rightarrow [T + U]_{\beta}^{\gamma} = [T]_{\beta}^{\gamma} + [U]_{\beta}^{\gamma}$$

**3. Let  $T: V_2(R) \rightarrow V_3(R)$  be the linear transformations defined by  $T(a_1, a_2) = (a_1 + 3a_2, 0, 2a_1 - 4a_2)$ . Let  $\beta$  and  $\gamma$  be the standard bases of  $V_2(R)$  and  $V_3(R)$  respectively. Verify  $[\alpha T]_{\beta}^{\gamma} = \alpha [T]_{\beta}^{\gamma}$**

Solution

$$\text{Given, } T(a_1, a_2) = (a_1 + 3a_2, 0, 2a_1 - 4a_2)$$

Since  $\beta$  and  $\gamma$  be the standard bases of  $V_2(R)$  and  $V_3(R)$

$$\text{the matrix corresponds to } \beta = [T]_{\beta}^{\gamma} = \begin{bmatrix} 1 & 3 \\ 0 & 0 \\ 2 & -4 \end{bmatrix}$$

the matrix corresponds to  $\beta = \alpha[T]_{\beta}^{\gamma} = \alpha \begin{bmatrix} 1 & 3 \\ 0 & 0 \\ 2 & -4 \end{bmatrix}$

$$= \begin{bmatrix} \alpha & 3\alpha \\ 0 & 0 \\ 2\alpha & -4\alpha \end{bmatrix} \dots (1)$$

We have,  $T(a_1, a_2) = (a_1 + 3a_2, 0, 2a_1 - 4a_2)$

$$\therefore \alpha T(a_1, a_2) = \alpha(a_1 + 3a_2, 0, 2a_1 - 4a_2)$$

$$(\alpha T)(a_1, a_2) = (\alpha a_1 + 3\alpha a_2, 0, 2\alpha a_1 - 4\alpha a_2)$$

Since  $\beta$  and  $\gamma$  be the standard bases of  $V_2(R)$  and  $V_3(R)$

the matrix corresponds to  $\beta = \alpha[T]_{\beta}^{\gamma} = \begin{bmatrix} \alpha & 3\alpha \\ 0 & 0 \\ 2\alpha & -4\alpha \end{bmatrix} \dots (1)$

$$\text{From (1) and (2)} \Rightarrow [\alpha T]_{\beta}^{\gamma} = \alpha [T]_{\beta}^{\gamma}$$

- 4. Let  $T: P_3(R) \rightarrow P_2(R)$  be the linear transformations defined by  $T(f(x)) = f'(x)$ . Let  $\beta$  and  $\gamma$  be the standard bases of  $P_3(R)$  and  $P_2(R)$  respectively. Then find  $[T]_{\beta}^{\gamma}$**

Solution :

Let,  $\beta = \{1, x, x^2, x^3\}$  be the standard bases of  $P_3(R)$

Let,  $\gamma = \{1, x, x^2\}$  be the standard bases of  $P_2(R)$

Let,  $w_1 = 1, w_2 = x, w_3 = x^2$

Given,  $T(f(x)) = f'(x)$ .

Let,  $(f(x)) = 1$ . Then  $f'(x) = 0$

$$\begin{aligned} T(1) &= T(f(x)) = f'(x) = 0 = 0.1 + 0.x + 0.x^2 \\ &= 0.w_1 + 0.w_2 + 0.w_3 \end{aligned}$$

The first column of  $[T]_{\beta}^{\gamma} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$

Let,  $(f(x)) = x$ . Then  $f'(x) = 1$

$$\begin{aligned} T(x) &= T(f(x)) = f'(x) = 1 = 1.1 + 0.x + 0.x^2 \\ &= 1.w_1 + 0.w_2 + 0.w_3 \end{aligned}$$

The second column of  $[T]_{\beta}^{\gamma} = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix}$

Let,  $(f(x)) = x^2$ . Then  $f'(x) = 2x$

$$\begin{aligned} T(x^2) &= T(f(x)) = f'(x) = 2x = 0.1 + 2.x + 0.x^2 \\ &= 0.w_1 + 2.w_2 + 0.w_3 \end{aligned}$$

The third column of  $[T]_{\beta}^{\gamma} = \begin{bmatrix} 0 \\ 2 \\ 0 \end{bmatrix}$

Let,  $(f(x)) = x^3$ . Then  $f'(x) = 3x^2$

$$\begin{aligned} T(x^3) &= T(f(x)) = f'(x) = 3x^2 = 0.1 + 0.x + 3.x^2 \\ &= 0.w_1 + 0.w_2 + 3.w_3 \end{aligned}$$

The fourth column of  $[T]_{\beta}^{\gamma} = \begin{bmatrix} 0 \\ 0 \\ 3 \end{bmatrix}$

$$\text{So, } [T]_{\beta}^{\gamma} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 2 & 0 \\ 0 & 0 & 0 & 3 \end{bmatrix}$$

